



Bloomington-Normal: 309-452-0995

Springfield: 217-717-4404

Satellite Offices:

Carlinville -Jacksonville - Morton

Sherman - Taylorville



Patient Name: _____ DOB: _____ MRN: _____

Epinephrine Auto-injector Authorization

To whom it may concern,

The patient named above is recommended to carry an epinephrine auto-injector, for the purpose of treating anaphylaxis, with them at all times. This medication is to be used in an emergency situation, along with other supportive monitoring and appropriate medications.

The patient will be carrying the following medication on her/his person:

- Epi Pen (0.3mg) or generic -- Mylan
- Epi Pen Jr. (0.15mg) or generic -- Mylan
- Auvi-Q (0.3mg)
- Auvi-Q 0.15 (0.15mg)
- Auvi-Q 0.1 (0.1mg)
- Adrenaclick (0.3mg) or generic -- Impax Labs
- Adrenaclick (0.15mg) or generic -- Impax Labs

Directions:

- Use as directed per the package insert, Intramuscular x1, into the upper outer thigh, perpendicular to the thigh.
- Hold the injector for 5-10 seconds and pull out the injector straight out to avoid skin laceration.
- Follow up immediately with a first-responder or health care professional.
- This injector may be repeated in 5-10 minutes if a second injector is available, should symptoms worsen or not improve, one time only.

Common Side Effects: increased heart rate, stronger or irregular heart rate, sweating, nausea and vomiting, difficulty breathing, paleness, dizziness, weakness or shakiness, headache, apprehension, nervousness or anxiety.

Please contact our office with any questions. You may email us as well: clinicalstaff@asthma2.com

Provider Name

Provider Signature

Date